



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# MULTIPLY PERFECT ODD NUMBERS WITH THREE PRIME FACTORS.

By R. D. CARMICHAEL.

The object of this note is to prove the following

PROPOSITION. There exist no multiply perfect odd numbers containing only three primes.

Let the numbers here considered be of the form  $p_1^{a_1} p_2^{a_2} p_3^{a_3}$ , where  $p_1, p_2, p_3$  are distinct odd primes, and  $p_1 < p_2 < p_3$ . And also let the multiplicity be  $m$ , where  $m > 1$ . Now, by definition, the multiplicity times the number equals the sum of the factors. Hence,

$$(1) \quad mp_1^{a_1} p_2^{a_2} p_3^{a_3} = \frac{p_1^{a_1+1}-1}{p_1-1} \cdot \frac{p_2^{a_2+1}-1}{p_2-1} \cdot \frac{p_3^{a_3+1}-1}{p_3-1}.$$

$$(2) \quad m = \frac{p_1^{a_1+1}-1}{p_1^{a_1}(p_1-1)} \cdot \frac{p_2^{a_2+1}-1}{p_2^{a_2}(p_2-1)} \cdot \frac{p_3^{a_3+1}-1}{p_3^{a_3}(p_3-1)}.$$

$$(3) \quad m < \frac{p_1}{p_1-1} \cdot \frac{p_2}{p_2-1} \cdot \frac{p_3}{p_3-1}.$$

The right member of (3) is greatest when the primes are smallest. By substituting 3, 5, 7 for  $p_1, p_2, p_3$ , we find that  $m \geq 2$ . Now, with  $m=2$  in (3), it is easily shown that we must have  $p_1=3, p_2=5, p_3 < 16$ , whence  $p_3=7, 11$ , or 13. Then (1) becomes

$$(4) \quad 2^4 \cdot 3^{a_1} \cdot 5^{a_2} \cdot p_3^{a_3} (p_3-1) = (3^{a_1+1}-1)(5^{a_2+1}-1)(p_3^{a_3+1}-1).$$

When  $p_3=7$ , equation (4) becomes

$$(5) \quad 2^5 \cdot 3^{a_1+1} \cdot 5^{a_2} \cdot 7^{a_3} = (3^{a_1+1}-1)(5^{a_2+1}-1)(7^{a_3+1}-1).$$

If  $a_3+1$  is even,  $7^{a_3+1}-1$  is divisible by  $7^2-1$ , which contains  $2^4$ . But, in any event,  $5^{a_2+1}-1$  contains  $2^2$  and  $3^{a_1+1}-1$  contains 2. The right member will then contain  $2^7$ , which is impossible. Hence,  $a_3+1$  is odd. Since odd powers of 7 end in 7 or 3,  $7^{a_3+1}-1$  is not divisible by 5. This requires  $a_1+1$  to be even; as odd powers of 3 end in 3 or 7, and  $3^{a_1+1}-1$  is therefore not then divisible by 5. Since  $a_1+1$  is even,  $3^{a_1+1}-1$  is divisible by  $3^2-1=2^3$ . The right member of (5) then contains  $2^6$ . This is impossible. Hence, there are no numbers of the type here considered when  $p_3=7$ .

Next, for  $p_3=11$ , (4) becomes

$$(6) \quad 2^5 \cdot 3^{a_1} \cdot 5^{a_2+1} \cdot 11^{a_3} = (3^{a_1+1}-1)(5^{a_2+1}-1)(11^{a_3+1}-1).$$

We can here, as above, show that  $a_3+1$  is odd. Likewise, that  $a_1+1$  is odd.

Then  $3^{a_1+1}-1$  does not contain 5. Hence, the right member contains the factor 5 only in  $11^{a_3+1}-1$ . But this factor must occur at least twice, as  $a_2 \neq 0$ . By writing  $(10+1)^{a_3+1}-1$  and expanding, we may easily show that it contains 5 only once unless  $a_3+1$  is divisible by 5. Then, let  $a_3+1=5n$ . Now,  $11^{5n}-1$  is divisible by  $11^5-1$ , which contains a prime greater than 11. Hence,  $p_3=11$  yields no numbers of the type here considered.

Finally, for  $p_3=13$ , (4) becomes

$$(7) \quad 2^6 \cdot 3^{a_1+1} \cdot 5^{a_2} \cdot 13^{a_3} = (3^{a_1+1}-1)(5^{a_2+1}-1)(13^{a_3+1}-1).$$

If  $a_3+1$  is even,  $13^{a_3+1}-1$  is divisible by  $13^2-1$ . This introduces the inadmissible factor 7. Hence,  $a_3+1$  is odd. The odd powers of 13 end in 3 or 7. Hence,  $13^{a_3+1}-1$  is not now divisible by 5. If  $a_1+1$  is odd,  $3^{a_1+1}-1$  is not divisible by 5. But to satisfy the equation, it must contain 5. Hence,  $a_1+1$  is even, and  $3^{a_1+1}-1$  then contains the factor  $3^2-1=2^3$ .  $5^{a_2+1}-1$  always contains  $2^2$ , and  $13^{a_3+1}-1$  always contains  $2^2$ . Hence, the right member contains  $2^7$ , which is impossible. Therefore, this case yields no numbers of the type here considered.

---

## DEPARTMENTS.

---

### ALGEBRA.

---

250. Proposed by PROFESSOR WILLIAM HOOVER, Ph. D., Athens, Ohio.

Factor  $a^2b^2(x^2+y^2)(a^2y^2+b^2x^2-a^2b^2)=(a^4y^2+b^4x^2)[1/(a^2y^2+b^2x^2)+ab]^2$ .

Solution by the PROPOSER.

Let  $x=r\cos\theta$ ..... (1),  $y=r\sin\theta$ ..... (2); then the given expression equated to zero becomes

$$\begin{aligned} (b^2\cos^2\theta+a^2\sin^2\theta)[a^2b^2-(b^4\cos^2\theta+a^4\cos^2\theta)]r^2 \\ -2ab\sqrt{[(b^2\cos^2\theta+a^2\sin^2\theta)](b^4\cos^2\theta+a^4\sin^2\theta)}r \\ =a^2b^2(b^4\cos^2\theta+a^4\sin^2\theta+a^2b^2) \dots\dots (3). \end{aligned}$$

Multiplying both sides of (3) by the coefficient of  $r^2$  and noticing that

$$\begin{aligned} a^2b^2-(b^4\cos^2\theta+a^4\sin^2\theta) &= a^2b^2(\sin^2\theta+\cos^2\theta)-(b^4\cos^2\theta+a^4\sin^2\theta) \\ &= b^2(a^2-b^2)\cos^2\theta+a^2(a^2-b^2)\sin^2\theta = (a^2-b^2)(b^2\cos^2\theta+a^2\sin^2\theta) \dots\dots (4), \end{aligned}$$

and similarly,

$$a^2b^2+b^4\cos^2\theta+a^4\sin^2\theta = (a^2+b^2)(b^2\cos^2\theta+a^2\sin^2\theta) \dots\dots (5).$$